arXiv:1406.4700v4 [math.GT] 17 Sep 2014

NON LEFT-ORDERABLE SURGERIES AND GENERALIZED BAUMSLAG-SOLITAR RELATORS

KAZUHIRO ICHIHARA AND YUKI TEMMA

Abstract. We show that a knot has a non left-orderable surgery if the knot group admits a generalized Baumslag-Solitar relator and satisfies certain conditions on a longitude of the knot. As an application, it is shown that certain positively twisted torus knots admit non left-orderable surgeries.

1. Introduction

In [3, Conjecture 1], Boyer, Gordon and Watson proposed the following conjecture, now called the *L-space Conjecture*: An irreducible rational homology 3-sphere is an L-space if and only if its fundamental group is non left-orderable. This has now become one of the important open problems in knot theory and Dehn surgery theory. Here, a rational homology sphere is called an *L-space* if the rank of the Heegaard Floer homology is equal to the order of the first (classical) homology group for the manifold. See [7]. Also a non-trivial group G is called *left-orderable*, often abbreviated as LO, if there exists a strict total ordering $>$ for the elements of G that is left-invariant: For any $f, g, h \in G$, whenever $g > h$ then $fg > fh$. We use the convention that the trivial group is not considered to be left-orderable.

On the study of the L-space Conjecture, one of the simple ways to construct L-spaces is given by Dehn surgery. For instance, any given knot in the 3-sphere S^3 admitting an L-space by surgery gives rise to an infinite family of L-spaces. In fact, it was shown in [8, Proposition 9.6] that if a knot K has one L-space surgery, then all the surgered manifolds are L-spaces if the surgery slopes are greater than or equal to $2g(K) - 1$, where $g(K)$ denotes the genus of the knot. Precisely, for a given knot K in a 3-manifold M, the following operation is called *Dehn surgery*; removing an open regular neighborhood of K from M, and gluing a solid torus back. Let us denote by $K(p/q)$ the 3-manifold obtained by Dehn surgery on K in S^3 along the slope r (i.e., the meridian of the attached solid torus is identified with the curve of the slope r), which corresponding to $p/q \in \mathbb{Q}$ on the peripheral torus of K. Please refer [2] on details, for instance.

We say that a Dehn surgery on a knot is *non left-orderable* if it yields a closed 3-manifold with the non left-orderable fundamental group. There are several known studies on 3-manifolds with a non left-orderable fundamental group by using Dehn surgery. For example, there are works by Clay and Watson [5] and Nakae [6]. In this paper, we show the following, which gives an extension of a result of Nakae.

Date: September 18, 2014.

2010 *Mathematics Subject Classification.* Primary 57M50; Secondary 57M25.

Key words and phrases. Dehn surgery, left-orderable group, Baumslag-Solitar relator, twisted torus knots.

Theorem 1.1. *Let K be a knot in a closed, connected 3-manifold M. Suppose that the knot group $\pi_1(M-K)$ has a presentation such as*

$$\langle a, b \mid (w_1 a^m w_1^{-1}) b^{-r} (w_2^{-1} a^n w_2) b^{r-k} \rangle$$

Here w_1, w_2 are arbitrary words with $m, n \geq 0$, $r \in \mathbb{Z}$, $k \geq 0$. Suppose further that a represents a meridian of K and $a^{-s} w a^{-t}$ represents a longitude of K with $s, t \in \mathbb{Z}$ and w is a word which excludes a^{-1} and b^{-1}. Then if $q \neq 0$ and $p/q \geq s+t$, then Dehn surgery on K along the slope p/q yields a closed 3-manifold with non left-orderable fundamental group.

There exist knots in S^3 whose knot groups admit presentations satisfying the conditions in our theorem. For instance, as shown by Nakae in [6, Proposition 2.1], the $(-2, 3, 2s+1)$-pretzel knot with $s \geq 3$ gives such examples. Thus our theorem presents an alternative proof of the result [6, Corollary 4.2].

Furthermore, in Proposition 3.1, we see that certain positively twisted torus knots have the knot groups admitting the presentations satisfying the conditions in our theorem. Therefore, as an application to Theorem 1.1, we show that certain positively twisted torus knots admit non left-orderable surgeries as follows.

Corollary 1.2. *Let K be a positively u-twisted $(3, 3v+2)$-torus knot with $u, v \geq 0$. Then if $q \neq 0$ and $p/q \geq 2u + 3(3v+2)$, then Dehn surgery on K along the slope p/q yields a closed 3-manifold with non left-orderable fundamental group.*

This gives an extension to the results obtained by Clay and Watson [5, Theorems 4.5 and 4.7]. There they gave proofs for the cases (1) $u \geq 0$ and $v = 1$, (2) $u = 1$ and $v \geq 0$. Also note that Vafaee showed in [10] that such twisted torus knots admit L-space surgery. Thus the corollary also gives a supporting evidence for the L-space conjecture.

We here remark that the relator in the presentation of $\pi_1(M-K)$ above can be regarded as a generalization of the well-known Baumslag-Solitar relator. By the *Baumslag-Solitar relator*, we mean the relator $x^{-n} y x^m y^{-1}$ with non-zero integers m and n in the group generated by two elements x and y. The groups with two generators and the Baumslag-Solitar relator was originally introduced in [1], now called a *Baumslag-Solitar group*, which plays an important role and is well-studied in combinatorial group theory and geometric group theory.

In particular, in [9], it was shown that the Baumslag-Solitar relator cannot appear in a non-degenerate way in the fundamental group of an orientable 3-manifold. Our relator is obtained from the Baumslag-Solitar relator by replacing x with some conjugates of it, and so it can be regarded as a generalization of the relator. It thus seems interesting that our relator can actually appear in the knot groups for certain knots in S^3, or more precisely, for the $(-2, 3, n)$-pretzel knots, and can play an essential role to study (non) left-orderability of the groups.

2. Proof

The following is the key proposition to prove the main theorem.

A homomorphism Φ of a group G to $Homeo^+(\mathbb{R})$, i.e., the group of order-preserving homeomorphisms on $\mathbb{R}$, has a *global fixed point* if there exists x in $\mathbb{R}$ such that $\Phi(g)x = x$ for all g in G. In the following, we will confuse, by abuse of notation g and $\Phi(g)$ for an element g in G.

Proposition 2.1. *Suppose that a group G has a presentation such as*

$$\langle a, b \mid (w_1 a^m w_1^{-1}) b^{-r} (w_2^{-1} a^n w_2) b^{r-k}, MLM^{-1}L^{-1}, M^p L^q \rangle$$

Here w_1, w_2 are arbitrary words with $m, n \geq 0$, $r \in \mathbb{Z}$, $k \geq 0$, $p, q \in \mathbb{Z}$, $M = a$, $L = a^{-s} w a^{-t}$, w is a word which contains at least one b and excludes a^{-1} and b^{-1}, $s, t \in \mathbb{Z}$. If $q \neq 0$ and $p/q \geq s + t$, then every homomorphism $\Phi : G \to Homeo^+(\mathbb{R})$ has a global fixed point.

Proof. We first consider the case where a has a fixed point, say x, on $\mathbb{R}$. In this case, we can show that $bx = x$ as follows. If b could not fix x, we can assume $bx > x$ for the x by choosing an order on $\mathbb{R}$. It is equivalent to $b^{-1}x < x$ since any action of G preserves an order of $\mathbb{R}$. Also $ax = x$, for x as above, is equivalent to $a^{-1}x = x$. Moreover, by the second relator $M^p L^q$ of G, it follows that

$$\begin{aligned}
M^p L^q &= a^p L^q \\
&= a^{p-(s+t)(q-1)-t} L a^{s+t} L a^{s+t} \cdots L a^{s+t} L a^t \\
&= a^{p-(s+t)(q-1)-t} (a^{-s} w a^{-t}) a^{s+t} (a^{-s} w a^{-t}) a^{s+t} \cdots (a^{-s} w a^{-t}) a^{s+t} (a^{-s} w a^{-t}) a^t \\
&= a^{p-(s+t)(q-1)-t-s} w^q \\
&= a^{p-(s+t)q} w^q \\
&= 1,
\end{aligned}$$

for $a = M$ and L commutes. Thus we have the relation $a^{p-(s+t)q} = w^{-q}$, and further that $a^{q(p-(s+t)q)} = w^{-q^2}$. On the other hand, we obtain $a^{-1}b^{-1}x < x$ by $a^{-1}b^{-1}x < a^{-1}x = x$. Repeating this consideration, we have $w^{-q^2}x < x$, for $q^2 > 0$ and w^{-1} is a word which contains at least one b^{-1} and excludes a and b. Therefore, we have

$$\begin{aligned}
x &> w^{-q^2} x \\
&= a^{q(p-(s+t)q)} x \\
&= x
\end{aligned}$$

giving a contradiction. Hence we have $bx = x$. Now x is fixed by a and b, and so $\Phi(G)$ has a global fixed point, since G is generated by a and b.

We next consider the case where a does not have any fixed point on $\mathbb{R}$. We will show that this case cannot happen. In this case, we can suppose that $ax > x$ for any $x \in \mathbb{R}$ by reversing its order if necessary. Using the first relator of G, we obtain the following:

$$\begin{aligned}
w_1 a^m w_1^{-1} b^{-r} w_2^{-1} a^n w_2 b^{r-k} &= 1 \\
w_1^{-1} b^{-r} w_2^{-1} a^n w_2 b^{r-k} w_1 a^m &= 1 \\
a^n w_2 b^{r-k} w_1 a^m &= w_2 b^r w_1
\end{aligned}$$

Then it follows that

$$\begin{aligned}
a^n w_2 b^{r-k} w_1 a^m x &= w_2 b^r w_1 x \\
&< w_2 b^r w_1 a^m x \\
&< a^n w_2 b^r w_1 a^m x
\end{aligned}$$

since we are assuming that $ax > x$ for any $x \in \mathbb{R}$. Since any element of G preserves an orientation of $\mathbb{R}$, we see that $a^n w_2 b^r w_1 a^m x > a^n w_2 b^{r-k} w_1 a^m x$ implies

$b^r w_1 a^m x > b^{r-k} w_1 a^m x$. Since the element $w_1 a^m \in G$ is thought as a homeomorphism of $\mathbb{R}$, there is a point $x \in \mathbb{R}$ which satisfies $x' = w_1 a^m x$ for any point $x' \in \mathbb{R}$. Hence, for the point x', we obtain

$$\begin{aligned} b^r x' &= b^r w_1 a^m x \\ &> b^{r-k} w_1 a^m x \\ &= b^{r-k} x' \\ b^k x' &> x' \end{aligned}$$

Consequently it follows that $bx > x$ for any $x \in \mathbb{R}$ if $k > 0$, and follows that $k \neq 0$ otherwise $x' > x'$.

On the other hand, by using the second relator of G, by similar arguments as above, it must follow that $a^{q((s+t)q-p)} x > x$ by $a^{q((s+t)q-p)} x = w^{q^2} x > x$. This must imply that $q((s+t)q-p) > 0$, that is, $p/q < s+t$, but this contradicts the assumption $p/q \geq s+t$.

This completes the proof of the proposition. □

Proof of Theorem 1.1. Suppose that $\pi_1(K(p/q))$ is left-orderable. It is known that a countable group G is left-orderable if and only if G is isomorphic with a subgroup of $Homeo^+(\mathbb{R})$. See [4, Theorem 2.6]. This implies that there exists an injective homomorphism $\Phi : \pi_1(K(p/q)) \to Homeo^+(\mathbb{R})$.

It was also proved in [4, Lemma 5.1] that if there is a homomorphism $G \to Homeo^+(\mathbb{R})$ with image $\neq \{id\}$, then there is another such homomorphism which induces an action on $\mathbb{R}$ without global fixed points. Without loss of generality, we may assume that Φ has no global fixed point. This contradicts the previous proposition. Therefore $\pi_1(K(p/q))$ is non left-orderable. □

3. Application

In this section, we show that the knot groups for certain twisted torus knots admit the presentations satisfying the conditions in our theorem, and give a proof of Corollary 1.2.

A *positively twisted torus knots* in S^3 is defined as the knots obtained from the torus knot by adding positive full twists along an adjacent pair of strands. Here we only consider positively u-twisted torus knots of type $(3, 3v+2)$, that is, the knots obtained from the torus knot of type $(3, 3v+2)$ by adding positive u full twists along an adjacent pair of strands. Note that it is known that the positively u-twisted $(3,5)$-torus knot is equivalent to the $(-2, 3, 5+2u)$-pretzel knot.

Proposition 3.1. *Let K be the positively u-twisted $(3, 3v+2)$-torus knot in S^3 with $u, v \geq 0$. Then the knot group $\pi_1(S^3 - K)$ has a presentation such as*

$$\langle a, b \mid (w_1 a^m w_1^{-1}) b^{-r} (w_2^{-1} a^n w_2) b^{r-k} \rangle$$

with $m = n = 1$, $r = u+1$, $k = 1$, $w_1 = (ba)^{v+1}$, $w_2 = (ba)^v$. Furthermore the generator a represents a meridian of K and the preferred longitude of K is represented as $a^{-s} w a^{-t}$ with $s = 2u + 3(3v+2) + 1$, $t = -1$ and $w = ((ba)^v b^{u+1})^2 (ba)^v b$.

Proof. As shown in [5, Proposition 3.1], $\pi_1(M - K)$ has the following presentation.

$$\langle x, y \mid x^2 (y^{-v} x)^u x y^{-v-1} (y^{-v} x)^{-u} y^{-2v-1} \rangle$$

Moreover by [5, Proposition 3.2], the element $y^{v+1}x^{-1}$ represents a meridian of K. Thus we set $a = y^{v+1}x^{-1}$. Then, since $a = y^{v+1}x^{-1}$ is equivalent to $x = a^{-1}y^{v+1}$, we have

$$\begin{aligned}&\langle x, y \mid x^2(y^{-v}x)^u xy^{-v-1}(y^{-v}x)^{-u}y^{-2v-1}\rangle\\ &= \langle a, y \mid (a^{-1}y^{v+1})^2(y^{-v}a^{-1}y^{v+1})^u a^{-1}y^{v+1}y^{-v-1}(y^{-v}a^{-1}y^{v+1})^{-u}y^{-2v-1}\rangle\end{aligned}$$

Let us transform the relator above as follows.

$$\begin{aligned}&(a^{-1}y^{v+1})^2(y^{-v}a^{-1}y^{v+1})^u a^{-1}y^{v+1}y^{-v-1}(y^{-v}a^{-1}y^{v+1})^{-u}y^{-2v-1}\\ &= (a^{-1}y^{v+1})(a^{-1}y^{v+1})(y^{-v}a^{-1}y^{v+1})^u a^{-1}(y^{-v}a^{-1}y^{v+1})^{-u}y^{-2v-1}\\ &= (a^{-1}y^{v+1})(a^{-1}y^{v+1})(y^{-v}a^{-1}y^{v+1})\cdots(y^{-v}a^{-1}y^{v+1})a^{-1}(y^{-v}a^{-1}y^{v+1})^{-u}y^{-2v-1}\\ &= (a^{-1}y^{v+1})a^{-1}ya^{-1}y\cdots ya^{-1}y^{v+1}a^{-1}(y^{-v}a^{-1}y^{v+1})^{-u}y^{-2v-1}\\ &= a^{-1}y^v(ya^{-1})^{u+1}y^{v+1}a^{-1}(y^{-v}a^{-1}y^{v+1})^{-u}y^{-2v-1}\\ &= a^{-1}y^v(ya^{-1})^{u+1}y^{v+1}a^{-1}(y^{-v-1}ay^v)^u y^{-2v-1}\\ &= a^{-1}y^v(ya^{-1})^{u+1}y^{v+1}a^{-1}(y^{-v-1}ay^v)\cdots(y^{-v-1}ay^v)y^{-2v-1}\\ &= a^{-1}y^v(ya^{-1})^{u+1}y^{v+1}a^{-1}y^{-v-1}ay^{-1}ay^{-1}\cdots ay^{-1}y^{-v}\\ &= a^{-1}y^v(ya^{-1})^{u+1}y^{v+1}a^{-1}y^{-v-1}(ay^{-1})^u y^{-v}\\ &= a^{-1}y^v(ya^{-1})^{u+1}y^{v+1}a^{-1}y^{-v-1}(ya^{-1})^{-u}y^{-v}\end{aligned}$$

Further, performing a cyclic permutation, we have the following.

$$(ya^{-1})^{-u}y^{-v}a^{-1}y^v(ya^{-1})^{u+1}y^{v+1}a^{-1}y^{-v-1}$$

Now we set $b = ya^{-1}$. Substituting $y = ba$, we have the following.

$$b^{-u}(ba)^{-v}a^{-1}(ba)^v b^{u+1}(ba)^{v+1}a^{-1}(ba)^{-v-1}$$

Finally, taking the inverse of the above, we have the following relator.

$$(ba)^{v+1}a(ba)^{-v-1}b^{-u-1}(ba)^{-v}a(ba)^v b^u$$

Consequently the knot group $\pi_1(S^3 - K)$ has a presentation such as

$$\langle a, b \mid (w_1 a^m w_1^{-1})b^{-r}(w_2^{-1}a^n w_2)b^{r-k}\rangle$$

with $m = n = 1$, $r = u + 1$, $k = 1$, $w_1 = (ba)^{v+1}$, $w_2 = (ba)^v$.

Also, by [5, Proposition 3.2], a longitude of K is represented by $x^2(y^{-v}x)^u x(y^{-v}x)^u$ in terms of the generators x and y. Then, by substituting $x = a^{-1}y^{v+1}$, it is transformed as follows.

$$\begin{aligned}&x^2(y^{-v}x)^u x(y^{-v}x)^u\\ &= (a^{-1}y^{v+1})^2(y^{-v}a^{-1}y^{v+1})^u a^{-1}y^{v+1}(y^{-v}a^{-1}y^{v+1})^u\\ &= (a^{-1}y^{v+1})(a^{-1}y^{v+1})(y^{-v}a^{-1}y^{v+1})\cdots(y^{-v}a^{-1}y^{v+1})a^{-1}y^{v+1}(y^{-v}a^{-1}y^{v+1})\cdots(y^{-v}a^{-1}y^{v+1})\\ &= (a^{-1}y^v)(ya^{-1})^{u+1}y^v(ya^{-1})^{u+1}y^{v+1}\end{aligned}$$

Again, by substituting $y = ba$, we obtain the following.

$$a^{-1}(ba)^v b^{u+1}(ba)^v b^{u+1}(ba)^{v+1}$$

Finally, as noted in the paragraph just below [5, Proposition 3.2], the presentation of the preferred longitude of K is obtained from the above by adding $a^{-3(3v+2)-2u}$ as follows.

$$a^{-3(3v+2)-2u-1}(((ba)^v b^{u+1})^2(ba)^v b)a$$

Therefore the preferred longitude of K is represented as $a^{-s}wa^{-t}$ with $s = 2u + 3(3v + 2) + 1$, $t = -1$ and $w = ((ba)^v b^{u+1})^2 (ba)^v b$.

This completes the proof of Proposition 3.1. □

Now Corollary 1.2 follows from Proposition 3.1 and Theorem 1.1 immediately.

Acknowledgements

The authors would like to thank Kimihiko Motegi for useful discussions. They also thank Anh Tran and Yasuharu Nakae for their valuable comments, and the anonymous referee for his/her careful readings for the the earlier version of this paper. The first author is partially supported by JSPS KAKENHI Grant Number 26400100.

References

[1] G. Baumslag and D. Solitar, Some two-generator one-relator non-Hopfian groups, *Bull. Amer. Math. Soc.* **68** (1962), 199–201.
[2] S. Boyer, Dehn surgery on knots, in *Handbook of geometric topology*, 165–218, North-Holland, Amsterdam.
[3] S. Boyer, C. McA. Gordon and L. Watson, On L-spaces and left-orderable fundamental groups, Math. Ann. **356** (2013), no. 4, 1213–1245.
[4] S. Boyer, D. Rolfsen and B. Wiest, Orderable 3-manifold groups, *Ann. Inst. Fourier (Grenoble)* **55** (2005), no. 1, 243–288.
[5] A. Clay and L. Watson, Left-orderable fundamental groups and Dehn surgery, *Int. Math. Res. Not.* **2013**, no. 12, 2862–2890.
[6] Y. Nakae, A good presentation of $(-2, 3, 2s+1)$-type pretzel knot group and $\mathbb{R}$-covered foliation, *J. Knot Theory Ramifications* **22** (2013), no. 1, 1250143, 23 pp.
[7] P. Ozsváth and Z. Szabó, On knot Floer homology and lens space surgeries, *Topology* **44** (2005), no. 6, 1281–1300.
[8] P. S. Ozsváth and Z. Szabó, Knot Floer homology and rational surgeries, *Algebr. Geom. Topol.* **11** (2011), no. 1, 1–68.
[9] P. B. Shalen, Three-manifolds and Baumslag-Solitar groups, *Topology Appl.* **110** (2001), no. 1, 113–118.
[10] F. Vafaee, On the Knot Floer Homology of Twisted Torus Knots, to apear in Int Math Res Notices (2014).

Department of Mathematics, College of Humanities and Sciences, Nihon University, 3-25-40 Sakurajosui, Setagaya-ku, Tokyo 156-8550, Japan
E-mail address: ichihara@math.chs.nihon-u.ac.jp

Graduate School of Integrated Basic Sciences, Nihon University, 3-25-40 Sakurajosui, Setagaya-ku, Tokyo 156-8550, Japan
E-mail address: s6113M12@math.chs.nihon-u.ac.jp